



Osmocote® Start

Start

Optimum results for salt-sensitive crops throughout the growing stage

11 | 11 | 17 | 2.0 | TE
N | P2O5 | K2O | MgO



Guaranteed analysis

oxide		
N	Total Nitrogen	11%
	Nitrate nitrogen (N-NO3)	4.5%
	Ammoniacal nitrogen (N-NH4)	6.5%
	Urea nitrogen (N-Urea)	0%
	Organic nitrogen	0%
P2O5	Phosphorus Pentoxide	11%
	Water soluble (P2O5)	8.2%
K2O	Potassium Oxide	17%
	Water Soluble (K2O)	17.0%
MgO	Magnesium Oxide	2.0%
	Water soluble (MgO)	1.0%
B	Boron	0.01%
	Water soluble (B)	0.01%
Cu	Copper	0.090%
	Water soluble (Cu)	0.080%
	Copper EDTA (Cu)	0.090%
Fe	Iron	0.38%
	Water soluble (Fe)	0%
	Iron EDTA (Fe)	0.17%
	Iron DTPA (Fe)	0%
	Iron EDDHA (Fe)	0%
Mn	Manganese	0.05%
	Water soluble (Mn)	0%
	Manganese EDTA (Mn)	0%
Mo	Molybdenum	0.009%
	Water soluble (Mo)	0.006%
Zn	Zinc	0.014%
	Water soluble (Zn)	0.007%
	Zinc EDTA (Zn)	0%

Description

Osmocote® Start coated fertiliser provides optimum results for short growth cycle plants. Perfect for violets, primulas, impatiens, marigolds, and many more. It is also ideal for salt-sensitive vegetable crops and for those cuttings proving difficult to root. As every single granule contains every element required, there is no risk of de-mixing any nutrients.

Benefits

- \\ Improve crop colour and root development
- \\ Increased efficiency with reduced leaching
- \\ Dust-free and easy to use

How to use

- 1 You can mix Osmocote® Start and use it for topdressing, just make sure not to leave any granules stuck to the leaves.
- 2 Store under dry conditions.
- 3 Properly seal partly used or damaged bags.
- 4 If you need more information, please contact your technical support.

Application rates

	Light feeding	Normal feeding
Container nursery stock	0.5-1.5 g/l	1.5-2.5 g/l
Pot plants	0.5-1.5 g/l	1.5-2.5 g/l
Bedding plants	0.5-2.0 g/l	-
Vegetable young plants	0.5-1.5 g/l	1.5-2.5 g/l

Attention These recommended rates are based on unfertilised substrates. Please be aware that these are general recommendations. Specific situations such as use in tunnels, greenhouses, or specific climate conditions require adjustments. This product is not recommended for dibbling. Contact your ICL Specialty Fertilisers advisor for more detailed advice. Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL Specialty Fertilisers cannot be held responsible for any adverse results.

Attention

Please contact your ICL Technical Area Sales Manager for more detailed advice.

ICL Growing Solutions marketing.ukire@icl-group.com

