



For the control of *Botrytis* (grey mould) on strawberries, raspberries, loganberries, blackberries and other *Rubus* hybrids, blackcurrants, redcurrants, whitecurrants, gooseberries and outdoor grapes.

MAPP 11229

Water dispersible granule formulation containing 50% w/w fenhexamid

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Bayer CropScience Limited
230 Cambridge Science Park
Milton Road,
Cambridge CB4 0WB
Telephone: 01223 226500

For 24 hour emergency information contact Bayer CropScience Limited

Telephone: 0800 220876

SAFETY PRECAUTIONS

Operator Protection

WASH CONCENTRATE from skin or eyes immediately
DO NOT BREATHE SPRAY
WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental Protection

Do not contaminate water with the product or its container.
Do not clean application equipment near surface water.
Avoid contamination via drains from farmyards and roads.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.
EMPTY CONTAINER COMPLETELY and dispose of safely.
Non-returnable containers must not be re-used for any purpose.

GB81732760b rA4a

PROTECT FROM FROST

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Teldor controls *Botrytis cinerea* (grey mould) on strawberries, cane fruit, bush fruit and outdoor grapes.

DISEASES CONTROLLED

For control of *Botrytis cinerea* (grey mould)

RESISTANCE MANAGEMENT

Alternating or mixing fungicides with different modes of action is a recognised disease resistance management strategy.

To minimise the likelihood of the development of *Botrytis* strains resistant to Teldor no more than 2 sprays of Teldor alone should be applied consecutively within a programme of sprays. Afterwards change to botryticides from different chemical groups for at least two consecutive sprays.

The possible development of disease strains resistant to Teldor cannot be excluded or predicted. Where such resistant strains occur, Teldor is unlikely to give satisfactory control.

CROP SPECIFIC INFORMATION

APPLICATION

For all crops listed on this label apply Teldor at 1.5 kg/ha.

Maximum individual dose: 1.5 kg/ha.

For successful control of *Botrytis* in all listed crops it is vital to achieve complete cover on all flowers and fruitlets throughout the blossom period. Sprayers should be set up in such a way as to achieve maximum cover on these parts, with sprays directed over the crop rows (not applied overall).

Do not use finer than 80 mesh filters.

Apply Teldor as a **MEDIUM** quality (BCPC definition) spray.

A maximum of 4 sprays of Teldor at 1.5 kg/ha may be applied in any season, however no more than 2 sprays of Teldor may be applied consecutively. Afterwards change to botryticides from a different chemical group for at least two consecutive sprays - see 'Diseases Controlled' above.

Strawberries

The first spray of a fungicide programme should be made at first flower (white bud).

Apply 1.5 kg Teldor in at least 300 litres of water per hectare, with a 10-14 day interval between sprays - see 'Number of Sprays' above.

Harvest Interval

At least 1 day must elapse between final treatment of Teldor and picking fruit.

Raspberries, Blackberries, Loganberries and other *Rubus* hybrid berries

The first spray of a fungicide programme should be made at early flower (white bud).

Apply 1.5 kg Teldor in at least 1,000 litres of water per hectare, with a 10-14 day interval between sprays - see 'Number of Sprays' above.

Harvest Interval

At least 1 day must elapse between final treatment of Teldor and picking fruit.

Currants, (black, red or white) and Gooseberries

The first spray of fungicide programme should be made at the late grape stage for currants and at first open flowers for gooseberries.

Apply 1.5 kg Teldor in at least 750 litres of water per hectare, with a 10-14 day interval between sprays - see 'Number of Sprays' above.

Harvest Interval

For currants (black, red and white) at least 3 days must elapse between final treatment of Teldor and picking fruit. For Goosberries at least 7 days must elapse between final treatment of Teldor and picking fruit.

Grapes - outdoor

Botrytis control usually requires a programme of fungicide sprays and Teldor may be used as part of such a programme.

If more than one application of Teldor at 1.5 kg/ha is made to a crop in the same growing season the applications must be alternated with a *Botrytis* control product from a different chemical group. If only two fungicide applications are made in a growing season for control of *Botrytis*, only one may include TELDOR - see 'Diseases Controlled' above.

Apply Teldor at 1.5 kg/ha as a protectant. Application may be made at any stage between full flower (BBCH 65) and start of berry softening (BBCH 85). Use a spray volume of 625 - 1,000 L/ha, depending on the application method, with a 21 day interval between sprays. Full coverage of flowers and developing fruit must be ensured by the application method employed.

Harvest Interval

At least 21 days must elapse between final treatment of Teldor and picking fruit.

MIXING

Half-fill the spray tank with clean water. Commence agitation. Steadily add the recommended quantity of Teldor directly into the water in the spray tank. Complete filling and maintain agitation of the suspension before and during spraying until the tank is empty.

® Teldor is a Registered Trade Mark of Bayer

© Bayer CropScience Limited 2015



To access the **Safety Data Sheet** for this product scan the code or use the link below:

www.bayercropscience.co.uk/teldorsds
or alternatively contact your supplier

Bayer

Bayer CropScience Limited 230 Cambridge Science Park Milton Road, Cambridge CB4 0WB Telephone: 01223 226500

Safety Information

TELDOR

Contains 50% w/w fenhexamid



Toxic to aquatic life with long lasting effects.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT INFORMATION

FOR USE ONLY AS A HORTICULTURAL FUNGICIDE

Crops: Strawberries, raspberries, loganberries, blackberries and other *Rubus* hybrids, blackcurrants, redcurrants, whitecurrants, gooseberries and outdoor grapes.

Maximum individual dose:

1.5 kg of product per hectare

Maximum number of treatments:

Two per year for grapes. Four per year for all other listed crops.

Latest time of application:

1 day before harvest for strawberries, raspberries, loganberries, blackberries and other *Rubus* hybrid berries.

3 days before harvest for currants (black, red or white) and 7 days before harvest for gooseberries.

21 days before harvest for outdoor grapes.

Other specific restrictions:

A maximum concentration of 0.24 kg of product in 100 litres of water must not be exceeded when making applications to outdoor grapes.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.